



Fig. 1A



Fig. 1B

PUROMYCIN-TETHERED OLIGO IS LIGATED TO mRNA (GENERATED FROM ABOVE CONSTRUCT) IN THE PRESENCE OF A SPLINT AND DNA LIGASE

NOTE: FOR SHORT ORF'S, THIS WHOLE TEMPLATE CAN BE MADE SYNTHETICALLY





Fig. 1C

IN VITRO TRANSLATION PROCEEDS NORMALLY FROM THE 5' TO THE 3' END OF THE mRNA

COVALENTLY LINKED PUROMYCIN ENTERS THE A SITE AND ATTACKS PEPTIDYL tRNA IN P SITE

RELEASE OF RNA-PROTEIN FUSION WITH HIGH SALT WASH OF RIBOSOME






Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6A




Fig. 6B




Fig. 6C




Fig. 6D




Fig. 6E




Fig. 6F




Fig. 6G




Fig. 6H




Fig. 7A




Fig. 7B




Fig. 7C




Fig. 8




Fig. 9

Replacement Sheet
 11/20




Fig. 10




Fig. 11




Fig. 12




Fig. 15A




Fig. 15B




Fig. 16A




Fig. 16B




Fig. 16C




Fig. 17




Fig. 13




Fig. 14




Fig. 18




Fig. 19




Fig. 20




Fig. 21




Fig. 22




Fig. 23




Fig. 24